

Circulating exosomal microRNAs as prognostic biomarkers in Cholangiocarcinoma: A Systematic Review

Gregorio Rangel^{1, 2, 3*}, Surasak Wanram⁴, Tsukuru Umemura⁵

¹Department of Biomedical Laboratory Science, Dili Institute of Health, Timor-Leste

²Department of Exacts and Natural Sciences, National Institute of Sciences and Technology, Timor-Leste

³Ossmali Research and Development, Timor-Leste

⁴College of Medicine and Public Health, Center for Excellence in Biomedical Science and Engineering, Ubon Ratchathani, 34190 Thailand

⁵Department of Medical Technology and Sciences, International University of Health and Welfare, Ohkawa, Fukuoka 831-8501, Japan

* Corresponding author Email: gregoriorangel20@gmail.com

Abstract

Introduction: Cholangiocarcinoma (CCA) is a cancer that arises from the cells within the bile ducts; both inside and outside the liver. Exosomal microRNAs (miRNAs) have been attracting major interest as potential biomarker in cancer. Our aims to do a systematic review on exosomal miRNAs serve as candidate clinical biomarkers in CCA. **Materials and methods:** Electronic databases (Medline, EMBASE, CINAHL and Cochrane data bases). We searched to identify publication all over the world related cancer and miRNA. PRISMA guidelines were followed. Selection was based on the design (Cholangiocarcinoma, invasive cholangiocarcinoma, intra-hepatic, extra-hepatic, liver cancer, microRNA, circulating miRNA, non-coding RNA, exosomal miRNA and exosomal vesicle), 27 papers were selected all over the world. CCA, target antigens, methodologies used for detection and miRNA expression were identified and summarized. **Results:** A total of 855 articles were searched, 333 duplicates removed and remain 522, 491 excluded after screened the titles and abstracts; and remain 31, 4 full texts articles were excluded with reason, and 27 identifications were included in the studies. Assay method was used to diagnose such as molecular Real Time-qPCR technique. **Conclusion:** The findings of this systematic review showed that microRNA functions as a tumor suppressor or a promoter in cholangiocarcinoma.

Keywords: Cholangiocarcinoma, intra-hepatic, extra-hepatic, microRNA, Medline, EMBASE, CINAHL, Cochrane, Systematic review.

1. Introduction

Cholangiocarcinoma (CCA) is an aggressive and fatal malignancy in the intra- and extra-hepatic biliary tract with increase in incidence and dismal prognosis (Wu, X, et al., 2016); according to the statistics, the incidence and mortality of CCA is raising worldwide (Nakanuma Y et al, 2015); this type of cancer frequently diagnosed as the second leading cause of cancer-related mortality worldwide (LV, H et al., 2018); MicroRNAs (miRNAs) are noncoding RNAs 18–25 nucleotides in length (Selaru, F. M, et al., 2009); the function of miRNA as posttranscriptional regulation of gene expression by either degradation of the targeting protein-coding RNAs or inhibition of their translation into protein (Wu, X, et al., 2016); the exosomal miRNAs in body fluids maybe useful diagnostic biomarker for the detection of the cancer (Ogata-kawata, H et al., 2014).

IFN is a family of cytokines critical not only for viral interference, but this cytokine can be inhibits cell proliferation and modulates differentiation,

apoptosis and migration (Fiorucci G, et al., 2014). An increasing quantity of researches concentration on microRNAs (miRNAs) which play multiple roles in variety of biological processes, including cancer (Huan, Q, et al., 2017). Most of CCA are unresectable when discovered since it has progressed into advanced stages. So the improvement in the diagnostic method of CCA is urgent, especially in biomarkers (Zhen, B, et al., 2017).

This literature review provides an overview of the most important scientific literature on CCA and exosomal miRNAs. Some scientific systematic reviews have been published in CCA such as cholecystectomy and the risk of cholangiocarcinoma, Non-alcoholic fatty liver disease as a risk factor for cholangiocarcinoma, preoperative biliary drainage in hilar cholangiocarcinoma, percutaneous vs. endoscopic pre-operative biliary drainage in hilar cholangiocarcinoma and identification of microRNAs as biomarkers for cholangiocarcinoma detection. However, there is no qualitative scientific systematic review of circulating exosomal miRNAs as biomarkers in cholangiocarcinoma. Therefore, reviewers concentrated on the scientific systematic

review on circulating exosomal miRNAs serves as candidate biomarker in cholangiocarcinoma.

2. Materials and Methods

Searching strategy

Reviewers are using the Cochrane guidelines to do a systematic review. The searching strategies used to conduct a systematic computerized search of the PubMed, Science Direct and Google Scholar databases (Bhagavathula, A. S., et al. 2017). However, in this review, reviewers accessed publication papers through Medline, EMBASE, and CINAHL to search the articles. Searching term that used as follows (i) "cholangiocarcinoma" or "intra-hepatic" or "extra-hepatic"; "liver cancer" (ii) "Circulating microRNA" or "Exosomal microRNA" or "miRNA" or "micro-RNA" or "short RNA" or "small RNA" or "non-coding RNA" (iii) "clinical trials" or "treatment" or "diagnosis" or "prognosis" or "recurrence" A detailed search strategy and search algorithms are shown in the searching results and summary tables.

Inclusion criteria

Research, review and systematic review articles were from trustworthy journals. These scientific articles involved studies reporting data from published papers including cholangiocarcinoma, intra-hepatic CCA, extra-hepatic CCA, liver cancer, pathobiology of biliary epithelial and human bile contain miRNAs extracellular vesicles. The inclusion criteria include: (1) studies must describe in human cholangiocarcinoma; (2) data on the intra-hepatic and extra-hepatic CCA; (3) data related exosomal microRNA and extracellular vesicle of cholangiocarcinoma; (4) data related liver cancer.

Exclusion criteria

In the exclusion, duplicate papers were removed, mismatch papers excluded based on the titles and abstract, and remain papers assessed for eligibility and included in the studies. Thus, the reviews exclude incredible publication papers on other types of cancer.

Review process

Research articles were identified from searches of the electronic databases was imported into ENDNOTE software version X8 (Thomson Reuters, USA). Before the data were extracted, selected articles were read the title and abstract to fulfill the inclusion criteria.

Data extraction and quality assessment

The inclusion and exclusion criteria were used to find the articles based on titles and abstracts. The selected articles were extracted and collected independently. The extraction data were included of the publication data (Authors, year of study, country of study, sample number, tumor stages, microRNA identified, follow-up months, detected samples, assay methods, normalizer RNA).

3. Results

The searching strategy was using Medline, EMBASE, CINAHL, and Cochrane. A total of 855 articles were searched, 333 duplicates removed and remain 522 articles, 491 excluded after screened the titles and abstracts, 4 full text articles were excluded with reason (the article study about long-non coding RNA). 27 articles were included in the studies (Figure 1).

A total of 30 articles, 18 were focused on CCA, 8 studies were related CCI, 1 study was concentrated on both ICC and ECC. Detected samples and tumor staging were assessed by molecular Real Time-qPCR techniques. List of miRNAs identified from the selected articles as shown in Figure 2 and summary description as indicated in Table 1.

4. Discussions

The miRNAs, a class of non-coding regulatory RNAs, have been detected in a variety of organism ranging from ancient unicellular eukaryotes to mammals. MicroRNAs (miRNAs) are non-coding RNAs of 20-22 nucleotides and regulate the translational inhibition of target mRNAs by base-pairing with their 3'-untranslated region (3'-UTR) (Li Jingjing et al., 2015). They have been associated with numerous molecular mechanisms involving developmental, physiological and pathological changes of cells and tissue (Judice, C. C., et al., 2016). Therefore, tissue, serum and plasma samples are required to investigate the existence of miRNAs. The miRNAs itself have been used as biomarkers to diagnose the infectious diseases, parasite, bacteria, including cancers. However, detection of miRNAs as biomarker in infectious diseases and cancers to evaluate the effectiveness of the drugs and diseases progression are still limited.

How microorganisms cause cancers

Helicobacter pylori are commonly transmitted person-to-person by saliva. The bacteria can also be spread by fecal contamination of food or water. Cheng-yeng kao et al., 2016 reported that after entering the host stomach, H. pylori utilize its urease activity to neutralize the hostile acidic condition at the beginning of infection. Schwabe, R. F. and C. Jobin., 2013 stated namely H. pylori infection substantially contributes to global cancer mortality. Meanwhile, Ringehan, M, et al., 2017 described that Hepatitis B virus (HBV) is one of the choric infection that leading cause for hepatocellular carcinoma (HCC) worldwide. Furthermore, Sithithaworn, P, et al., 2014 defined explicitly that liver fluke infection causes pathological changes mainly to the bile ducts where the worm can be found, as well as to the liver and gall bladder in both human and animal. The early pathological changes consisted of an acute inflammatory reaction involving the large intrahepatic bile ducts and portal connective tissue. Therefore, the infection

of the bacteria, viruses and parasites may produce specific miRNAs during the infection process that leading to cancers.

MicroRNA in cancer

In human, microorganism (Bacteria, virus, parasites) invasion associated gene expression and immune system which is related with diseases severity and assessed drug effectiveness. Li Pang et al., 2018 declared that for the expression, miRNAs are first transcribed by RNA polymerase II and processed to form hairpin-like intermediates called pre-miRNAs. Bohnsack MT et al., 2004 reported Exportin-5 then transports these pre-miRNAs out of the nucleus and into the cytoplasm for further processing, converting the pre-miRNA into conserved. Bartel, D. P., 2009 defined It is these mature miRNAs of 18–25 nucleotides in length that can control cellular function through recognition of specific targets by complementary base pairing, forming either RNA-induced silencing complexes (RISC), inducing mRNA degradation or inhibiting mRNA translation endogenous, small noncoding mature RNAs as showed in the figure 2.

Gabay M, et al., 2014 defined MYC is a transcription factor that modulates the gene expression of thousands of genes that regulate many programs which are hallmarks of cancer including: metabolism, proliferation, self-renewal, and survival. Dhanasekaran R, et al., 2017 defined MYC's ability to maintain proliferation, survival and self-renewal were regulated via its induction of miR17 ~ 92 clusters. Rangel, G., et al., 2019 stated candidate miRNAs could be investigated using 3 different algorithms that are the most widely used in the updated version as follows: miRanda, RegRNA and Target Scan. Kanaoka R., et al., 2017 showed recent studies have indicated that miRNAs can be detected in biological fluids such as plasma and serum and hold promise as noninvasive biomarkers for cancer patients. Jiao, D, et al, 2017 described the differential miRNA expression patterns were validated with the TaqMan qRT-PCR assay.

Candidate miRNAs in Cholangiocarcinoma

The miRNAs are a group of small non-coding RNAs, which can bind to mRNAs of target gene leading to their degradation or translational inhibition. Chiang, N. J, et al 2015 defined that the roles of miRNA in the biology of CCA, such as proliferation, invasion, migration, differentiation and apoptosis. Rizvi S and Gores G, J., 2017 reported that CCAs are epithelial tumors with markers of cholangiocyte differentiation, anatomically CCAs are classified into intrahepatic (iCCA), perihilar (pCCA) and distal (dCCA) subtypes. Dutta S et al., 2015 indicated that exosomes are small (40–100 nm in diameter) membrane bound vesicles that are initially formed within the endosomal compartment and secreted upon fusion of the limiting membrane of multi-vesicular bodies (MVBs) with plasma membrane. Kongpetch et al., 2015 identified that miRNA in CCA cell lines

(HUCCT1 and MEC) revealed biliary epithelial cell-specific miRNAs, i.e., miR22, miR125a, miR127, miR199a, miR199a, miR214, miR376a and miR424, which are down-regulated in these lines. Selaru et al., 2009 detected in a separate study, miR21 was found to be up regulated in ICC compared to normal epithelial bile duct tissue. Inhibition of miR21 was shown to increase protein expression of PDCD4 and TIMP3 which are the inhibitors of program-cell death and metastasis, respectively.

Candidates' genes that associated with miRNAs such as FOXA1 associated with miR-212 reported by Zhou J, et al., 2017; Zbtb7a associated with miR-106a informed by Jiao et al., 2017; miR-21 associated with PDCD4 as reported by Selaru et al., 2009. The mentioned genes have studied and showed that FOXA1 was correlated with miR-212 in the intrahepatic cholangiocarcinoma; Zbtb7a was linked with miR-106a which the studied was focused on miRNA expression and sensitivity of CCA cell. However, in this scientific systematic review, reviewers will study the candidates' miRNAs in both intrahepatic CCA and extra-hepatic CCA in the future perspective.

A 24 selected articles were included in the studies, microorganism infection, cell proliferation, migration and differentiation might be produced specific miRNAs which associated with PDCD4 and other genes such as iRBE, RBE, CEA, pCEA, TET1-p53, Rac1 were showed that the majority of genes that associated with miRNAs (miR-21, miR-221, miR-26-5p, miR-451, miR-150, miR-191 and miR-122) were detected during the infection, cell proliferation, cell differentiation and promoted tumor invasion and metastasis. These miR-150, miR-191, miR-26, miR-21, miR-451 and miR-122 were associated with ICC samples after detected by molecular PCR. The miR-21 also identified from ECC sample. These miRNAs would be selected as candidates of miRNAs as biomarker of human CCA in the future perspective.

MicroRNA 21

There were 6 studies assessing miR-21 to detect this miRNA expression and its regulates and promoting cell proliferation, cell development and metastasis in CCA (Gallego et al., 2016, Garajova et al., 2015, Huang, Q, et al., 2013, Liu, N, et al 2015, Selaru et al., 2009, Wang, L, et al.,2015). Gallego et al. used 12 human tissue samples to compare with 40 normal plasma specimens in RT-qPCR. The result indicated that there were 4 miRNAs (miR-21, miR-34c, miR-200b, and miR-221) highly up-regulated in the tumor samples. These miRNAs recommended for further analysis in samples. Garajova et al. isolated 69 samples from ICC and ECC. Real Time PCR was used to investigate the specimens. The result has shown that miR-21 expression as prognostic biomarker. Therefore, further investigation of ICC and ECC in plasma. Qiu Huang et al. identified miR-21 was significantly higher perineural invasion and lymph node metastasis. The samples were used qRT-PCR

for studied. Liu et al. stated namely CCA tumor cell line was isolated and cultured. RNA was reverse transcribed into cDNA using Primer Script RT reagent kit (Takara, China). The result presented miR-21 regulates biological behavior by inducing EMT in human CCA. Selaru et al. argued that 20 primary CCA were compared with 14 normal tissues, using qRT-PCR detect miR-21 was shown highly expressed. Wang, L, et al. reported that expression of miR-21 in culture media ICC lines and serum plasma compared with healthy control subjects, using RT-PCR identify miR-21 was shown promoted ICC proliferation and cell growth in vitro by targeting PTPN14 and PTEN.

MicroRNA-150

There are 3 studies that assessed miR-150 in CCA to detect the role of the miRNA itself as tumor suppressor gene involved tumor invasion and metastasis and regulates cell proliferation and cell differentiation (Fan F et al., 2016, Wang et al., 2014, Wu et al., 2016). An F et al. reported that miR-150-5p continuously decreased in the serum of sclerosing cholangitis and body fluids of CCA. Microarray was used to analyze the serum samples. The results showed that over expression of miR-150 attenuated the growth, invasion and migration capability of CCA cells with target oncogene Ets like gene-1 (ELK1). Wang et al. described expression miR-150 in tissues and plasma sample from patients compared with normal tissues using quantitative reverse transcription polymerase chain reaction (qRT-PCR) indicated miR-150 patient's plasma sample was up-regulated with noncancerous plasma. Wu et al. stated 28 CCAs and 30 samples from primary sclerosing cholangitis (PSCs) were compared with 50 healthy controls. Serum, bile and tissues samples were used to diagnose in qRT-PCR. The result presented that miR-150 lower expression if compared normal plasma controls.

MicroRNA 191 and miR-451

A study (Li et al., 2017) described namely 84 samples from ICC evaluated by RT-PCR indicated that miR-191 up-regulated expressed from ICC tissues compared with adjacent bile duct tissues. Thus, miR-191 also promoted cholangiocarcinoma cell Tsukuru Umemura, Professor of the University

proliferation in human. On one hand, (LV et al., 2014) reported 26 clinical tissue patients were investigated by RT-PCR. The result designated that miR-451 under-expressed compared to the peritumoral tissues.

5. Conclusion and Prospective

MicroRNAs are small non-coding RNAs that regulate gene expression. They exert their effects on the cells they are synthesized in, and are also released into the extracellular space and transported in body fluids such as blood and urine. Exosomal miRNAs may have important functions in cell-cell communication and have potential as biomarkers to detect and monitor disease. Limited prognosis in CCA as one of the most lethal malignant disease. Thus, low study cover on miRNA in CCA itself. Therefore, reviewers decide 4 candidate miRNAs such as miR-21, miR-150, miR-191 and miR-451 as exosomal circulating miRNAs. Further investigation of the plasma, serum and tissues samples in the laboratory needed performed using real time PCR towards selected candidate miRNAs especially for ICC and ECC samples in the future perspective.

6. Additional Information and Declarations

Funding

The study was funded by Dili Health Science Institute
Grant Disclosures

The following grant information was disclosed by the authors:

Dili Health Science Institute, Timor-Leste (Fund number)

Completing Interests

The authors declare there are no competing interests.

Author Contributions

Gregorio Rangel, wrote the papers.

Surasak Wanram, Advisor and Final editing of the paper




Figure 1. The searching strategy in the systematic review, searching papers using MEDLINE, EMBASE, CINAHL and Cochrane, duplicates papers were removed. The papers screened based on the titles and abstracts and irrelevant excluded. Relevant papers assessed for eligibility (Some irrelevant articles excluded with reason) included articles in the studies.

Biogenesis of miRNAs




Figure 2 Biogenesis of miRNA. (1) DNA transcription and non-coding genome region. (2) Primary-miRNA (Pri-miRNA) transcribed by RNA polymerase II. (3) Drosha enzyme cut precursor miRNA (Pre-miRNA) and pre-miRNA transferred out of nucleus into cytoplasm with exporting 5. (4) Dicer cut pre-miRNA convert become duplex intermediate. (5) Duplex transform developed into mature miRNA. (6) MiRNAs degradation. (7) Infected cells and immune system produce cytokines in early infection, severe and cerebral malaria. (8) MiRNAs suppress the expression of the target genes via mRNA cleavage or translation repression. The functions of miRNAs are involved in inflammation, epigenetic change, metastasis, and apoptosis of the cell to promote to CCA cells.

Selected References

- Bernuzzi F, Marabita F, Lleo A, Carbone M, Miolo M, Marzoni M, et al. Serum microRNAs as novel biomarkers for primary sclerosing cholangitis and cholangiocarcinoma. *Clinical & Experimental Immunology*. 2016;185(1):61-71.
- Chen L, Yan H-X, Yang W, Hu L, Yu L-X, Liu Q, et al. The role of microRNA expression pattern in human intrahepatic cholangiocarcinoma. *Journal of hepatology*. 2009;50(2):358-69.
- Chen Y, Liu D, Liu P, Chen Y, Yu H, Zhang Q. Identification of biomarkers of intrahepatic cholangiocarcinoma via integrated analysis of mRNA and miRNA microarray data. *Molecular medicine reports*. 2017;15(3):1051-6.
- Cheng Q, Feng F, Zhu L, Zheng Y, Luo X, Liu C, et al. Circulating miR-106a is a novel prognostic and lymph node metastasis indicator for cholangiocarcinoma. *Scientific reports*. 2015; 5:16103.
- Chusom P, Namwat N, Loilome W, Techasen A, Pairojkul C, Khuntikeo N, et al. Overexpression of microRNA-21 regulating PDCD4 during tumorigenesis of liver fluke-associated cholangiocarcinoma contributes to tumor growth

- and metastasis. *Tumor Biology*. 2013;34(3):1579-88.
- Correa-Gallego C, Maddalo D, Doussot A, Kemeny N, Kingham TP, Allen PJ, et al. Circulating plasma levels of microRNA-21 and microRNA-221 are potential diagnostic markers for primary intrahepatic cholangiocarcinoma. *PloS one*. 2016;11(9):e0163699.
- Fan F, Lu J, Yu W, Zhang Y, Xu S, Pang L, et al. MicroRNA-26b-5p regulates cell proliferation, invasion and metastasis in human intrahepatic cholangiocarcinoma by targeting S100A7. *Oncology Letters*. 2018;15(1):386-92.
- Huang Q, Liu L, Liu C-H, You H, Shao F, Xie F, et al. MicroRNA-21 regulates the invasion and metastasis in cholangiocarcinoma and may be a potential biomarker for cancer prognosis. *Asian Pacific Journal of Cancer Prevention*. 2013;14(2):829-34.
- Kawahigashi Y, Mishima T, Mizuguchi Y, Arima Y, Yokomuro S, Kanda T, et al. MicroRNA profiling of human intrahepatic cholangiocarcinoma cell lines reveals biliary epithelial cell-specific microRNAs. *Journal of Nippon Medical School*. 2009;76(4):188-97.
- Kim NH, Kim H, Park SK, Yang HJ, Jung YS, Park JH, et al. Sa1802 Decreased miRNA-145-5p Mediated Upregulation of Rac1 Differentially Stimulates Cholangiocarcinoma Cell Growth, Migration and Invasion. *Gastroenterology*. 2016;150(4):S370.
- Kwon H, Song K, Han C, Zhang J, Lu L, Chen W, et al. Epigenetic Silencing of miRNA-34a in Human Cholangiocarcinoma via EZH2 and DNA Methylation: Impact on Regulation of Notch Pathway. *The American journal of pathology*. 2017;187(10):2288-99.
- Li H, Zhou ZQ, Yang ZR, Tong DN, Guan J, Shi BJ, et al. MicroRNA-191 acts as a tumor promoter by modulating the TET1-p53 pathway in intrahepatic cholangiocarcinoma. *Hepatology*. 2017;66(1):136-51.
- Li J, Gao B, Huang Z, Duan T, Li D, Zhang S, et al. Prognostic significance of microRNA-203 in cholangiocarcinoma. *International journal of clinical and experimental pathology*. 2015;8(8):9512.
- Li Z, Shen J, Chan MT, Wu WKK. The role of microRNAs in intrahepatic cholangiocarcinoma. *Journal of cellular and molecular medicine*. 2017;21(1):177-84.
- Liang Z, Liu X, Zhang Q, Wang C, Zhao Y. Diagnostic value of microRNAs as biomarkers for cholangiocarcinoma. *Digestive and Liver Disease*. 2016;48(10):1227-32.
- Liu N, Jiang F, He T-L, Zhang J-K, Zhao J, Wang C, et al. The roles of MicroRNA-122 overexpression in inhibiting proliferation and invasion and stimulating apoptosis of human cholangiocarcinoma cells. *Scientific reports*. 2015; 5:16566.
- Liu Z, Jin Z-Y, Liu C-H, Xie F, Lin X-S, Huang Q. MicroRNA-21 regulates biological

behavior by inducing EMT in human cholangiocarcinoma. *International journal of clinical and experimental pathology*. 2015;8(5):4684.

18. Piontek K, Selaru FM, editors. *MicroRNAs in the biology and diagnosis of cholangiocarcinoma*. *Seminars in liver disease*; 2015: Thieme Medical Publishers.

19. Razumilava N, Bronk SF, Smoot RL, Fingas CD, Werneburg NW, Roberts LR, et al. miR-25 targets TNF-related apoptosis inducing ligand (TRAIL) death receptor-4 and promotes apoptosis resistance in cholangiocarcinoma. *Hepatology*. 2012;55(2):465-75.

20. Selaru FM, Oлару AV, Kan T, David S, Cheng Y, Mori Y, et al. MicroRNA-21 is overexpressed in human cholangiocarcinoma and regulates programmed cell death 4 and tissue inhibitor of metalloproteinase 3. *Hepatology*. 2009;49(5):1595-601.

21. Stutes M, Tran S, DeMorrow S. Genetic and epigenetic changes associated with cholangiocarcinoma: from DNA methylation to microRNAs. *World journal of gastroenterology: WJG*. 2007;13(48):6465.

22. Wang L-J, He C-C, Sui X, Cai M-J, Zhou C-Y, Ma J-L, et al. MiR-21 promotes intrahepatic cholangiocarcinoma proliferation and growth in vitro and in vivo by targeting PTPN14 and PTEN. *Oncotarget*. 2015;6(8):5932.

23. Wang S, Yin J, Li T, Yuan L, Wang D, He J, et al. Upregulated circulating miR-150 is associated with the risk of intrahepatic cholangiocarcinoma. *Oncology reports*. 2015;33(2):819-25.

24. Zhang M-Y, Li S-H, Huang G-L, Lin G-H, Shuang Z-Y, Lao X-M, et al. Identification of a novel microRNA signature associated with intrahepatic cholangiocarcinoma (ICC) patient prognosis. *BMC cancer*. 2015;15(1):64.

Table 1 Summary table of selected papers include in the systematic review

No	Identification	Year of publication	Country of study	Clinical samples	Biomarker category	Target genes	Tumor type	Assay method	MicroRNAs identified	Regulation	Control
1	F. Bernuzzi et al	2016	UK	Serum	Diagnostic	CEA	PSC and CCA	qRT-PCR	miR-222, miR-483-5p, miR-194	Up	Healthy
2	L. Chen et al	2009	China	Serum	Diagnostic	pCEA	ICC	qRT-PCR	miR-21, miR-25, miR-223 and miR-222 and miR-198	Up and Down	Healthy
3	Chen qiabao et al	2015	China	Serum	Prognostic	CEA	CCA	qRT-PCR	miR-106a	Down	Healthy
4	P. Chusorn et al	2013	Thailand	Cell line	Diagnostic	PDCD4	ICC	qRT-PCR	miR-21	Up	GAPDH
5	Correa-Gallego et al	2016	USA	Plasma	Diagnostic	PDCD4	ICC	qRT-PCR	miR-21, miR-34, miR-200b, miR-221	Up	Healthy
6	Fan et al	2017	China	Tissue	Diagnostic	iRBE	ICC	qRT-PCR	miR-26b-5p	Down	Healthy
7	Qian Huan et al	2013	China	Tissue	Diagnostic	RBE	CCA	qRT-PCR	miR-21	Up	U6
8	Kawahigashi et al	2009	Jepang	Cell line	Diagnostic	HuCC1, MEC, TFK-1, IHGGK	ICC	qRT-PCR	miR-22, miR-127, miR-125a	Down	U6
9	Kim et al	2016	South Korea	SNU-245	Diagnostic	SRGAP2, Rac1.	ECC	qRT-PCR	miR-145-5p	Up	Normal
10	Kwon et al	2017	South Korea	Serum	Diagnostic	Notch1, Notch2, Jagged1	CCA	qRT-PCR	miR-34a	Down	GAPDH
11	Li Hao et al	2017	China	Tissue	Diagnostic	TET1-p53	ICC	qRT-PCR	miR-191	Up	GAPDH
12	Li Jingjing et al	2015	China	Tissue	Prognostic, Diagnostic	miR-203	CCA	qRT-PCR	miR-203	Up	U6
13	Lin zheng et al	2016	China	NP	NP	NP	ICC	NP	NP	NP	NP
14	Liang et al	2016	China	Tissue	Prognostic	NP	CCA	Pubmed and Embasse	NP	NP	NP
15	Chen yaqing et al	2017	China	Tissue	Prognostic	TIAM1	ICC	Microarray data	miR-21	NP	NP
16	Liu Ning et al	2015	China	Tissue	Diagnostic	RBE	CCA	qRT-PCR	miR-122	Down	Normal
17	Lin zheng et al	2016	China	Cell line	Diagnostic	RBE	CCA	qRT-PCR	miR-21	Down	GAPDH
18	Piontek and Selaru	2015	USA	NP	NP	CCA cells	CCA	NP	NP	NP	NP
19	Razumilava et al	2012	USA	Cell line	Diagnostic	MCM7	CCA	qRT-PCR	miR-25, miR-106b	Up	Normal
20	Selaru et al	2009	USA	Tissue	Diagnostic	PDCD4	CCA	qRT-PCR	miR-21	Up	U6
21	Stutes et al	2009	USA	NP	NP	CCA cells	CCA	NP	NP	NP	NP
22	Wang li-juang et al	2015	China	Serum, cell line	Diagnostic	RBE, HUCCT1	ICC	qRT-PCR	miR-21	Up	U6
23	Wang shouli et al	2014	China	Tissue	Diagnostic	CEA	ICC	qRT-PCR	miR-150, miR-638	Down	Negative
24	Zhang et al	2015	China	Tissue	Diagnostic	CEA	ICC	qRT-PCR	miR-612, miR-105-5p	Down, Up	U6

Remark: CEA = Carcinoembryonic antigen, PSC = primary sclerosing cholangitis, CTR = Control, PDCD = Programmed Cell Death, GAPDH = Glyceraldehyde-3-phosphate dehydrogenase, FNH = focal nodular hyperplasia

References

1. Wu, X., et al. (2016). "Profiling of downregulated blood-circulating miR-150-5p as a novel tumor marker for cholangiocarcinoma." *Tumor Biology* 37(11): 15019-15029.
2. Nakanuma, Y. and Y. Kakuda (2015). "Pathologic classification of cholangiocarcinoma: new concepts." *Best Practice & Research Clinical Gastroenterology* 29(2): 277-293.
3. Lv, H., et al. (2018). "Noncoding RNAs in liver cancer stem cells: the big impact of little things." *Cancer letters* 418: 51-63.
4. Selaru, F. M., et al. (2009). "MicroRNA-21 is overexpressed in human cholangiocarcinoma and regulates programmed cell death 4 and tissue inhibitor of metalloproteinase 3." *Hepatology* 49(5): 1595-1601.
5. Ogata-Kawata, H., et al. (2014). "Circulating exosomal microRNAs as biomarkers of colon cancer." *PloS one* 9(4).
6. Fiorucci, G., et al. (2015). "MicroRNAs in virus-induced tumorigenesis and IFN system." *Cytokine & growth factor reviews* 26(2): 183-194.
7. Huang, Q., et al. (2017). "Circulating microRNAs in colorectal cancer." *JMolGenetMed* 11 274: 1747-0862.1000274.
8. Zheng, B., et al. (2017). "miRNA and lncRNA as biomarkers in cholangiocarcinoma (CCA)." *Oncotarget* 8(59): 100819.
9. Bhagavathula, A. S., et al. (2017). "Anti-malarial treatment outcomes in Ethiopia: a systematic review and meta-analysis." *Malaria journal* 16(1): 269.
10. Li, J., et al. (2015). "Prognostic significance of microRNA-203 in cholangiocarcinoma." *International journal of clinical and experimental pathology* 8(8): 9512.
11. Judice, C. C., et al. (2016). "MicroRNAs in the host-apicomplexan parasites interactions: a review of immunopathological aspects." *Frontiers in cellular and infection microbiology* 6: 5.
12. Chen, Y., et al. (2017). "Identification of biomarkers of intrahepatic cholangiocarcinoma via integrated analysis of mRNA and miRNA microarray data." *Molecular medicine reports* 15(3): 1051-1056.
13. Schwabe, R. F. and C. Jobin (2013). "The microbiome and cancer." *Nature Reviews Cancer* 13(11): 800-812.
14. Ringehan, M., et al. (2017). "Viral hepatitis and liver cancer." *Philosophical Transactions of the Royal Society B: Biological Sciences* 372(1732): 20160274.
15. Sithithaworn, P., et al. (2014). Roles of liver fluke infection as risk factor for cholangiocarcinoma. *J Hepatobiliary Pancreat Sci*, 21, 301-8.
16. Li, P., et al. (2018). "MicroRNA-192-5p is a predictive biomarker of survival for Stage IIIB colon cancer patients." *Japanese journal of clinical oncology* 48(7): 619-624.
17. Bohnsack, M. T., et al. (2004). "Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs." *Rna* 10(2): 185-191.
18. Bartel, D. P. (2009). "MicroRNAs: target recognition and regulatory functions." *cell* 136(2): 215-233.
19. Gabay, M., et al. (2014). "MYC activation is a hallmark of cancer initiation and maintenance." *Cold Spring Harbor perspectives in medicine* 4(6): a014241.
20. Dhanasekaran, R., et al. (2018). "Anti-miR-17 therapy delays tumorigenesis in MYC-driven hepatocellular carcinoma (HCC)." *Oncotarget* 9(5): 5517.
21. Rangel, G., et al. (2020). "Candidate microRNAs as Biomarkers in Malaria Infection: A Systematic Review." *Current Molecular Medicine* 20(1): 36-43.
22. Kanaoka, R., et al. (2018). "Usefulness of plasma exosomal microRNA-451a as a noninvasive biomarker for early prediction of recurrence and prognosis of non-small cell lung cancer." *Oncology* 94(5): 311-323.
23. Jiao, D., et al. (2017). "miR-106b regulates the 5-fluorouracil resistance by targeting Zbtb7a in cholangiocarcinoma." *Oncotarget* 8(32): 52913.
24. Chiang, N.-J., et al. (2015). "Epigenetic regulation in the carcinogenesis of cholangiocarcinoma." *The international journal of biochemistry & cell biology* 67: 110-114.
25. Rizvi, S. and G. J. Gores (2017). "Emerging molecular therapeutic targets for cholangiocarcinoma." *Journal of hepatology* 67(3): 632-644.
26. Dutta, S., et al. (2015). "Proteomics profiling of cholangiocarcinoma exosomes: a potential role of oncogenic protein transferring in cancer progression." *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease* 1852(9): 1989-1999.
27. Kongpetch, S., et al. (2015). "Pathogenesis of cholangiocarcinoma: From genetics to signalling pathways." *Best Practice & Research Clinical Gastroenterology* 29(2): 233-244.
28. Zhu, L., et al. (2017). "[Retracted] MicroRNA-212 targets FOXA1 and suppresses the proliferation and invasion of intrahepatic cholangiocarcinoma cells." *Experimental and therapeutic medicine* 13(5): 2109-2109.
29. Correa-Gallego, C., et al. (2016). "Circulating plasma levels of microRNA-21 and microRNA-221 are potential diagnostic markers for primary intrahepatic cholangiocarcinoma." *PloS one* 11(9).
30. Garajová, I., et al. (2015). "MiR-21 expression as prognostic biomarker in extrahepatic but not intrahepatic radically resected cholangiocarcinomas." *Annals of Oncology* 26: vi98.
31. Huang, Q., et al. (2013). "MicroRNA-21 regulates the invasion and metastasis in cholangiocarcinoma and may be a potential biomarker for cancer prognosis." *Asian Pacific Journal of Cancer Prevention* 14(2): 829-834.
32. Liu, N., et al. (2015). "The roles of MicroRNA-122 overexpression in inhibiting proliferation and invasion and stimulating apoptosis of human

cholangiocarcinoma cells." *Scientific reports* 5: 16566.

33. Wang, L.-J., et al. (2015). "MiR-21 promotes intrahepatic cholangiocarcinoma proliferation and growth in vitro and in vivo by targeting PTPN14 and PTEN." *Oncotarget* 6(8): 5932.

34. Fan, F., et al. (2018). "MicroRNA-26b-5p regulates cell proliferation, invasion and metastasis in human intrahepatic cholangiocarcinoma by targeting S100A7." *Oncology Letters* 15(1): 386-392.

35. Li, H., et al. (2017). "MicroRNA-191 acts as a tumor promoter by modulating the TET1-p53 pathway in intrahepatic cholangiocarcinoma." *Hepatology* 66(1): 136-151.

Author Profile



Biography: Gregorio Rangel is one civil servant of National Health Laboratory, Ministry of Health, Timor-Leste. He was completed his post graduate diploma in applied parasitology and entomology at institute for medical research (IMR), Kuala Lumpur, Malaysia in 2014. He was concluded his master's degree in biomedical science in Ubon Ratchathani University, Thailand in 2018. In government institutions, He was work with National Malaria Program, Ministry of Health from 2009-2013, and National Health Laboratory from 2013-2021. In partnership development, He has been worked as national consultant for World Health Organization Timor-Leste Gregorio Rangel from November – December, 2020 with project "Infection Prevention and Control". He has also been worked as national consultant for World Health Organization Timor-Leste from September 2021-July 2022, project entitled "Essential Health Services during COVID-19 period". He has been also worked as national assessor for National Agency for Academic Assessment and Accreditation (NAAAA) from July-September 2022. He is a young scientist in medical laboratory. He has contributed various research and publication in biomedical science. Currently, he is working as head of department (equivalent national director) of exacts and natural science, National Institute for science and technology, Timor-Leste. He also a full-time lecturer at department of Biomedical Laboratory Science, Dili Health Science Institute, Timor-Leste. Meanwhile, he is working as principal investigator in a collaborative study between Menzies School of Health Research in Timor-Leste, Ministry of Health and Dili Health Science Institute with project entitled "The Use of Highly Sensitive Diagnostic and Typing/Sequencing Technology in the Pursuit of Malaria Elimination in Timor Leste".